

The Alexander polynomial: the algebraic way

①

Aarnout: If $G \cong \frac{F(x_1, \dots, x_n)}{\langle r_1, \dots, r_s \rangle}$, consider the maps

$$\frac{\partial}{\partial x_i} : \mathbb{Z}[F_n] \rightarrow \mathbb{Z}[F_n]$$

$$\mathbb{Z}[F_n] \xrightarrow{\gamma} \mathbb{Z}[G] \xrightarrow{\alpha} \mathbb{Z}[G^{ab}]$$

ρ

where $F_n = F(x_1, \dots, x_n)$ free group in n letters $x_1, \dots, x_n$,

γ induced by the projection $F_n \rightarrow F_n / \langle r_i \rangle \cong G$

α induced by the abelianization $G \rightarrow G / [G, G] = G^{ab}$.

Then the Alexander matrix of $\langle x_i \mid r_j \rangle$ is $A = \left(\rho \left(\frac{\partial r_j}{\partial x_i} \right) \right)$.

Theorem: The elementary ideals $E_k(A)$ do not depend on the choice of presentation of G .

Aim: Create a new Knot invariant out of a presentation of the knot group




$\pi_K := \pi_1(S^3 - K)$ using the elementary ideals of the Alexander matrix of π_K .

Since π_K is already a Knot invariant, what we create out of this will also be a Knot invariant.

Proposition: let K be a knot. Then $\pi_K^{ab} \cong \mathbb{Z}$.

Pf: By the Wirtinger algorithm, $\pi_K \cong \frac{F(x_1, \dots, x_n)}{\langle r_1, \dots, r_n \rangle}$,

where r_i is one of the following two relations:



$$r_i: x_k x_i = x_{i+1} x_k$$

$$x_k x_{i+1} = x_i x_k$$

In the abelianization, the relation r_i becomes $x_i = x_{i+1}$, so

$$\left(\frac{F(x_1, \dots, x_n)}{\langle r_1, \dots, r_n \rangle} \right)^{ab} \cong F(x_1) \cong \mathbb{Z} = \langle t \rangle$$

$x_i \longleftrightarrow x_1 \longleftrightarrow 1 \longleftrightarrow t$

□

Exercise: Show that if L is a link of l components, then $\pi_1(S^3 - L)^{ab} \cong \mathbb{Z}^l$.

The upshot is that our group ring at hand is $\mathbb{Z}[\langle t \rangle] = \mathbb{Z}[t, t^{-1}]$, the ring of Laurent polynomials. Any element is of the form

$$p(t) = \sum_{i=n}^m a_i t^i, \quad n \leq m, \quad n, m \in \mathbb{Z}, \quad a_i \in \mathbb{Z}.$$

(2)

Lemma: The ring $\mathbb{Z}[t, t^{-1}]$ is a unique factorization domain, and its invertible elements are $\mathbb{Z}[t, t^{-1}]^\times = \{\pm t^{\pm n}\}$.

Pf: $\mathbb{Z}$ is a UFD (Euclidean), therefore so is $\mathbb{Z}[t]$ (Gauss), and therefore

$$\mathbb{Z}[t][t^{-1}] = S^{-1}\mathbb{Z}[t], \text{ where } S = \{1, t, t^2, \dots\} \text{ (the localization of } \mathbb{Z}[t] \text{ at } S)$$

UFD is a UFD). $\square$

For the second part, we let $\varphi: \mathbb{Z}[t, t^{-1}] \rightarrow \mathbb{Z}[t]$ be defined by

$$\varphi\left(p(t) = \sum_{i=-n}^m a_i t^i\right) := t^{-n} \cdot p(t)$$

It is routine to check that φ is a ring homomorphism. Then, if $p(t) \in \mathbb{Z}[t, t^{-1}]^\times$ with inverse $q(t)$, we have

$$1 = \varphi(1) = \varphi(p(t) \cdot q(t)) = \varphi(p(t)) \cdot \varphi(q(t)).$$

Since the only units in $\mathbb{Z}[t]$ are ± 1 , we get that $\varphi(p(t)) = \pm 1$ and

$$\text{therefore } p(t) = \pm t^{-n}.$$

 $\square$

* In a unique factorization domain, we can define the gcd of a finite number of elements (and the proof is the same as for $\mathbb{Z}$).

• To sum up, given a knot K we can consider a collection of ideals $E_r(A) \subset \mathbb{Z}[t, t^{-1}]$, where A is the Alexander matrix of π_K .

Definition: let K be a knot. The r -th Alexander polynomial $\Delta_r(t)$ of K is the generator of the smallest principal ideal $(\Delta_r(t)) \subset \mathbb{Z}[t, t^{-1}]$ containing $E_r(A)$, that is,

$$\Delta_r(t) := \gcd(E_r(A)).$$

The generator is unique up to an invertible element of $\mathbb{Z}[t, t^{-1}]$, ie, up to multiplication by $\pm t^{\pm n}$.

The first Alexander polynomial $\Delta_1(t)$ is called the Alexander polynomial of K and is denoted by $\Delta_K(t)$. In other words,

$$\Delta_K(t) = \gcd((n-1) \times (n-1) \text{ minors of } A).$$

• We can make the computation a bit easier:

Fact: In the Wirtinger presentation $\pi_K \cong \langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ of a knot K , one of the relations r_i (any) is superfluous, that is, it is a consequence of the rest of relations.

(3)

- Consequently, π_K admits a presentation $\langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$, so a Alexander matrix of dimension $(n-1) \times n$.

Corollary: $\Sigma_0(A) = 0$, so $\Delta_0(t) = 0$.

- At this point we would still have to compute n determinants to get $\Delta_K(t)$. The following proposition asserts that it is enough to take one!

Proposition: let A be the Alexander matrix of a knot $A \in \mathcal{M}_{n \times n}(\mathbb{Z}[t, t^{-1}])$ obtained from a presentation of π_K given by the Wirtinger algorithm. Then A is equivalent to any of its $(n-1) \times (n-1)$ submatrices. Consequently, if A' is one of them then

$$\Delta_K(t) = \det A'.$$

Pf: We already saw that we can drop one of the rows of A . Let us see now that we can drop any column. The key point is that, as in calculus, the free calculus satisfies the equality

$$f(x) - f(1) = \sum_{j=1}^n \frac{\partial f}{\partial x_j} \cdot (x_j - 1)$$

(here $f(1)$ stands for the image of f under the morphism $\mathbb{Z}[F_n] \rightarrow \mathbb{Z}$, i.e.,

$$x_i \mapsto 1$$

$f(1)$ is obtained by substituting every x_i by 1). Applying this to $f = r_i$ we get

$$r_i - 1 = \sum_{j=1}^n \frac{\partial r_i}{\partial x_j} (x_j - 1).$$

Since $\rho(r_i) = 1$ and $\rho(x_k) = \rho(x_j) = t \quad \forall k, j = 1, \dots, n$,
 applying ρ to the latter expression we get

$$\begin{aligned} 0 &= \rho(r_i) - 1 = \sum_{j=1}^n \rho\left(\frac{\partial r_i}{\partial x_j}\right) \cdot (\rho(x_j) - 1) \\ &= (t-1) \cdot \sum_{j=1}^n a_{ij} \end{aligned}$$

Since $(t-1) \neq 0$ and $\mathbb{Z}[t, t^{-1}]$ is a integral domain, $\sum_{j=1}^n a_{ij} = 0$, i.e., any column is linear combination of the rest. $\square$

Remark: Since we already know the expression of any r_i , it provides us a quick way to compute $\Delta_k(t)$: namely, for a negative crossing, r_i is $x_k x_i = x_{i+1} x_k$.

The only non-zero derivatives are

$$\frac{\partial r_i}{\partial x_k} = 1 - x_i, \quad \frac{\partial r_i}{\partial x_i} = x_k, \quad \frac{\partial r_i}{\partial x_{i+1}} = -1$$

After abelianizing, this becomes


$$a_{ik} = 1-t, \quad a_{ii} = t, \quad a_{i, i+1} = -1.$$

For a positive crossing, the coefficients are

$$a_{ik} = 1-t, \quad a_{ii} = -1, \quad a_{i, i+1} = t.$$

Later we only have to take the determinant of any $(n-1) \times (n-1)$ submatrix. (4)

Example: Figure-eight knot 4_1



$$A = \begin{matrix} & \begin{matrix} x_1 & x_2 & x_3 & x_4 \end{matrix} \\ \begin{matrix} r_1 \\ r_2 \\ r_3 \\ r_4 \end{matrix} & \begin{pmatrix} t & -1 & 0 & 1-t \\ 1-t & -1 & t & 0 \\ 0 & 1-t & t & -1 \\ t & 0 & 1-t & -1 \end{pmatrix} \end{matrix}$$

$$\Delta_K(t) = \det (\text{any } 3 \times 3 \text{ submatrix})$$

$$= \begin{vmatrix} t & -1 & 0 \\ 1-t & -1 & t \\ 0 & 1-t & t \end{vmatrix} = -t^2 - t^2(1-t) + t(1-t)$$

$$= -t^2 - t^2 + t^3 + t - t^2 = t^3 - 3t^2 + t = \underline{\underline{t^2 - 3t + 1}}$$

⑩

Let us find the value of the determinant of the matrix

formed by the elements of the matrix

$$A = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$



(continued from page 3) $\Delta = 11, \Delta$

$$(1+1) + (1+1) - 1 = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix}$$

$$1 + 18 - 1 = 1 + 18 - 1 = 18$$